


PRVPATENT- OCH REGISTRERINGSVERKET
Patentavdelningen**BEST AVAILABLE COPY****Intyg
Certificate**

Härmed intygas att bifogade kopior överensstämmer med de handlingar som ursprungligen ingivits till Patent- och registreringsverket i nedannämnda ansökan.

This is to certify that the annexed is a true copy of the documents as originally filed with the Patent- and Registration Office in connection with the following patent application.



(71) Sökande Akzo Nobel nv, Arnhem NL
Applicant (s)

(21) Patentansökningsnummer 0302986-5
Patent application number

(86) Ingivningsdatum 2003-11-13
Date of filing

EPO - DG 1

22. 11. 2004

Stockholm, 2004-11-10

(71)

För Patent- och registreringsverket
For the Patent- and Registration Office.

Hjärdís Segerlund

Avgift
Fee 170:-

**PRIORITY
DOCUMENT**SUBMITTED OR TRANSMITTED IN
COMPLIANCE WITH RULE 17.1(a) OR (b)

USE OF A DERIVATIVE OF ASPARTIC ACID AS A COLLECTOR IN FROTH FLOTATION PROCESSES

5 The present invention relates to the use of a derivative
of aspartic acid as a collector for a phosphate containing
mineral, such as apatite, in a froth flotation process.
According to the invention the collector has a high
selectivity for phosphate containing minerals even in the
presence of carbonate minerals, such as calcite. A method for
10 the manufacture of the derivatives is also disclosed.

Phosphate rocks contain calcium phosphate minerals
largely in the form of apatite usually together with other
minerals for example silicate minerals and carbonate
minerals, such as calcite. Apatite is a generic name for a
15 group of calcium phosphate minerals also containing other
elements or radicals such as fluorapatite, chlorapatite,
carbonate apatite and hydroxyl apatite.

It is well-known to separate the valuable phosphate
minerals from the barren minerals by using a froth flotation
20 process where the phosphate minerals are enriched in the
float. In these flotation processes fatty acids and naphthenic
acids and their soaps have frequently been used as a
collector. However, this type of collectors works well only
when silicate minerals are the barren mineral. When carbonate
25 minerals, such as calcite, are present in the ore, a low
selectivity for the phosphate minerals is obtained. The
selectivity can to a certain degree be improved by the
concurrent use of depressants, such as polysaccharides of
different types.

30 Anionic surfactants such as alkylbenzene sulphonates,
alkyl phosphates and alkyl sulphosuccinamates have also been
proposed as flotation agents for phosphate containing ores,


but their selectivity for and yield of calcium phosphate in froth flotation processes are still too low.

In US patent 4 358 368 it is disclosed that the selectivity for calcium phosphate minerals can be essentially improved by using amphoteric surfactants of the sarcosinate type. The sarcosinate is advantageously used in combination with a nonionic, water-insoluble polar co-collector. The drawback of sarcosinate as a collector is the fact that it has limited ability to fasten to the apatite surfaces which limits the yield of apatite in the concentrate.

Further, the US Patent 4 043 902 discloses a process for froth flotation of non-sulfide ores such as sulfates, carbonates, fluorides, tungstates, phosphates and oxides, e.g. celestite, barite, sheelite, fluorite, calcite, magnesite, gypsum, anhydrite, cassiterite, apatite and the like, using salts of tri- and tetra-carboxyl containing fatty alkyl substituted aspartic acids, aspartic mono-esters, and aspartic di-esters, as collectors in conjunction with appropriate gangue depressants where required.

The US Patent 4 790 932 describes a process for the froth flotation of non-sulfidic mineral containing ores, in which process an anionic and/or nonionic collector surfactant is used as a collector in conjunction with at least one N-alkyl or N-alkenyl aspartic acid as a co-collector.

According to the present invention it has now been found that a certain derivative of aspartic acid has excellent properties as a collector for a calcium phosphate-containing mineral in an alkaline froth flotation process of an ore also containing calcium carbonate. The derivative of the invention has the formula



(I),


where R^I is a hydrophobic group containing a hydrocarbon group of 6-24 carbon atoms; R^{II} is an alkyl group with 1-7 carbon atoms, preferably 1-3 carbon atoms, or a group of the formula $(B)_yH$, in which B is an alkyleneoxy group with 2-4 carbon atoms and y is a number from 1 to 10, preferably from 1 to 3; and M is a group selected from the group consisting of a cation or hydrogen. Preferably R^I is a glycidylether group of the formula $CH_2CH(OH)CH_2O(A_1)_{x1}R_1$, in which R_1 is a hydrocarbon group with 8-24 carbon atoms, A_1 is an alkyleneoxy group with 2-4 carbon atoms and $x1$ is a number from 0 to 10, preferably from 0 to 5; a hydroxyl group of the formula $CH_2CH(OH)R_2$, in which R_2 is a hydrocarbon group with 6-22 carbon atoms; a propylene ether group of the formula $C_3H_6O(A_3)_{x3}R_3$, in which R_3 is a hydrocarbon group with 8-24 carbon atoms, A_3 is an alkyleneoxy group with 2-4 carbon atoms and $x3$ is a number from 0-10, preferably from 0 to 5; or of the formula R_4 , where R_4 is a hydrocarbon group containing 8-24 carbon atoms. Suitably the group $(A_1)_{x1}R_1$ is $(C_2H_4O)_{1-3}R_1$, where R_1 is a hydrocarbon group of 10-20 carbon atoms, such as an aliphatic group or an alkylphenyl group, while $x3$ is 0-3. Most preferably R^{II} is methyl, hydroxyethyl or hydroxypropyl. The cation M is normally a monovalent cation, such as sodium, potassium or an ammonium cation. The amount of the derivative can vary within wide limits but is normally between 10 and 1500, preferably between 50 and 800, grams per ton of the ore.

The froth flotation process of the invention results in a high concentration and a high yield of calcium phosphates in the float. The derivatives of the invention are suitably used in combination with a nonionic, water-insoluble polar compound as a co-collector, whereby the selectivity and the yield is further improved. The polar co-collector has a good


affinity for the particles coated with the derivative and can thereby improve or further enhance the properties of the derivative. The co-collector can be used in amounts between 0 and 1000, preferably between 5 and 350, grams per ton of the ore.

The derivative of the invention can be manufactured by reaction steps well-known to a person skilled in the art. For example, under alkaline conditions, maleic acid or a salt thereof can be reacted with

- 10 a) a primary amine of the formula $R^{II}NH_2$, where R^{II} has the meaning mentioned above, followed by reacting the intermediate obtained with a glycidylether of the formula




where R_1 , $x1$ and A_1 have the meanings mentioned above, an epoxide of the formula




where R_2 has the meaning mentioned above, or a halide compound of the formula $HalR_4$, where Hal is a halide and R_4 has the meaning above; or

- 25 b) with a primary amine of the formula $R^I NH_2$, where R^I has the meaning mentioned above, followed by reacting the intermediate obtained with a halide compound of the formula $HalR^{II}$, where Hal is a halide and R^{II} has the meaning mentioned above.


30 A more specific method of producing the derivative according to the invention is to react for example the disodium salt of maleic acid with methylamine to obtain the N-methylaspartic acid disodium salt. This reaction product can then be further reacted with a compound




where R_1 , A_1 and $x1$ have the meanings mentioned above, to
 5 obtain an aspartate of the formula



Another method is to react the intermediate product, N-methylaspartate disodium salt, with a compound of the formula




where R_2 has the meaning mentioned above, to an aspartate of
 15 the formula




Still another method is to first react the monosodium salt of maleic acid with a compound of the formula


25 $R_3(A_3)_{x3}\text{OC}_3\text{H}_6\text{NH}_2$, where R_3 , A_3 and $x3$ have the meanings mentioned above, to obtain an aspartate intermediate of the formula



The intermediate can then be reacted with $\text{ClCH}_2\text{CH}_2\text{OH}$ or CH_3Cl and NaOH to form a derivative of the formulae



and




The polar co-collector is suitably an alkylene oxide adduct of the formula



in which R^{III} is a hydrocarbon group, preferably an aliphatic group or an alkylphenyl group, with 8-22 carbon atoms, A is an oxyalkylene group having 2-4 carbon atoms and p is a number from 1-6. The oxyalkylene groups are suitably oxyethylene groups or a mixture of oxyethylene and oxypropylene groups. By placing the oxypropylene groups and especially the oxybutylene groups in the end position of the adduct, a lower foaming is achieved.

Another suitable co-collector is an ester of the formula



in which R^{IV} is an aliphatic group having 7-21 carbon atoms, A is an alkyleneoxy group having 2-4 carbon atoms, q is a number from 0-6, and Y is an alkyl group having 1-4 carbon atoms or hydrogen, provided that Y cannot be hydrogen when q is zero.

In addition to their advantageous froth flotation effect, the co-collectors also have a favourable effect on foaming by making the foam less stable when used in combination with the derivative of the invention.

In the process according to the invention, it is also possible to add pH-adjusting substances, such as sodium carbonate and sodium hydroxide, foaming agents, foam regulators, depressants, such as waterglass, different types of starch and CMC, and activating substances. In the present

froth flotation process the pH-value of the pulp is suitably within the range of 8-11.

The present invention is further illustrated by the following working examples.

5

Example 1

A magmatic ore, containing about 12% by weight of fluorapatite and about 73% by weight of calcite and a rest containing silicates and magnetite, was ground to a particle size of $\leq 630 \mu\text{m}$. The ground ore in an amount of 390 grams, 10 0.8 liter of water and 78 mg of hydrolysed corn starch dissolved in an amount of 1% by weight in water, were added to a flotation cell of 1.5 liter, whereupon the pH value was adjusted to 10.5 by addition of NaOH and the ground ore was conditioned for 5 minutes at 23°C. After the conditioning, 78 15 mg of a reagent according to the table below was added as a 1% by weight solution in water and the total amount in the flotation cell was adjusted by addition of water to 1.4 liter. The content of the flotation cell was then conditioned for 2 minutes, followed by a rougher flotation step and one 20 or more cleaning steps of the rougher concentrates.

The rougher concentrate and the concentrates from the cleaning steps were analysed with regard to their contents of phosphate (P_2O_5) and calcite. The results obtained are shown in Table II below.

11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
66
67
68
69
70
71
72
73
74
75
76
77
78
79
80
81
82
83
84
85
86
87
88
89
90
91
92
93
94
95
96
97
98
99
100

Table I. Reagents

Code		Composition
A	a)	39% by weight of sarcosinate of the formula $\text{nonylphenyl}-(\text{C}_2\text{H}_4\text{O})_{1,1}\text{CH}_2\text{CH}(\text{OH})\text{CH}_2\text{N}^+(\text{CH}_3)\text{CH}_2\text{COO}^-$ <p>according to the US patent 4 358 368</p>
	b)	27% by weight of the reaction product between 1 mole of nonylphenol and 2 moles of ethylene oxide
	c)	34% by weight of a solvent consisting of water and propylene glycol
B	a)	39% by weight of $\text{nonylphenyl}-(\text{C}_2\text{H}_4\text{O})_{1,1}\text{CH}_2\text{CH}(\text{OH})\text{CH}_2\text{NHCH}(\text{CH}_2\text{COONa})\text{COONa}$
	b)	As in reagent A, b)
	c)	As in reagent A, c)
1	a)	39% by weight of $\text{nonylphenyl}-(\text{C}_2\text{H}_4\text{O})_{1,1}\text{CH}_2\text{CH}(\text{OH})\text{CH}_2\text{N}(\text{CH}_3)\text{CH}(\text{CH}_2\text{COONa})\text{COONa}$
	b)	As in reagent A, b)
	c)	As in reagent A, c)
2	a)	39% by weight of $\text{nonylphenyl}-(\text{OC}_2\text{H}_4)_{1,1}\text{OC}_3\text{H}_6\text{-N}(\text{C}_2\text{H}_4\text{OH})\text{CH}(\text{CH}_2\text{COONa})\text{COONa}$
	b)	As in reagent A above
	c)	As in reagent A above

The reagent A represents the prior art and B is a comparison, while the aspartate-containing reagents 1 and 2 are in accordance with the invention.

5

Table II. Flotation results of reagents A, 1 and 2

Reagent	Flotation step	Concentrate		
		Content, % P_2O_5	Yield, % apatite	Content, % calcite
A	Rougher	18.3	99.0	43.2
	Cleaning 1	38.8	68.0	2.3
B	Rougher	19.9	93.6	-
	Cleaning 1	29.5	86.2	-
	Cleaning 2	36.7	77.9	-
	Cleaning 3	41.7	64.2	-
1	Rougher	17.3	98.5	43.2
	Cleaning 1	31.9	96.5	9.1
	Cleaning 2	40.2	92.3	2.0
	Cleaning 3	42.5	87.3	0.5
2	Rougher	23.2	94.3	41.4
	Cleaning 1	34.0	90.8	15.9
	Cleaning 2	38.7	85.4	4.5
	Cleaning 3	40.2	78.3	2.0

The results show that the aspartate-containing reagents are superior to the reagent A in accordance with the prior art and the comparison B. The content and yield of apatite are improved, while the content of calcite is low.

Example 2

500 g of a magnetic ore having a particle size of $\leq 5 \mu m$ and containing 9% by weight of fluorapatite, 17% by weight of calcite and a rest mainly consisting of silicates was ground

in a rod mill together with 0.4 liter of water, 180 mg of NaOH and 50 mg of waterglass with a ratio between SiO₂ and Na₂O of 3.3:1 to a particle size, where 80% by weight of the ground ore had a particle size ≤250 μm. The ground ore, 125 mg of the reagent in Table 3, and water were added to a flotation cell of 1.5 liter, the water being added in such an amount that the total volume of the ore pulp became 1.4 liter. After adjusting the pH value to 11 by the addition of NaOH, the pulp was conditioned at 21°C for 5 minutes. To the conditioned pulp, 25 mg of an iso-butyric acid ester of secondary butanol was added as a foamer, and a rougher flotation step was performed followed by three cleaning steps. The concentrates from the rougher flotation and from the cleaning steps were analysed with regards to the yield of apatite and the results obtained are shown in the Table IV below.

Table III. Reagent

Code		Composition
3	a)	36% by weight of an aspartate of the formula $ \begin{array}{c} \text{CH}_3 \\ \\ \text{R}-\text{CH}_2\text{CHCH}_2\text{N}-\text{CH}-\text{CH}_2\text{COONa}, \\ \qquad \qquad \\ \text{OH} \qquad \qquad \text{CH}_2\text{COONa} \end{array} $ <p>where R is an aliphatic group containing 13-15 carbon atoms</p>
	b)	21% by weight of the reaction product between 1 mole of nonylphenol and 2 moles of ethylene oxide
	c)	43% by weight of a solvent consisting of water and propylene glycol

Table IV. Flotation results of reagent 3

Reagent	Flotation step	Concentrate	
		Content, % P_2O_5	Yield, % apatite
3	Rougher	18.5	97.1
	Cleaning 1	23.7	94.8
	Cleaning 2	32.1	90.6
	Cleaning 3	37.5	74.5

The results show that the reagent according to the invention makes it possible to increase the content of apatite and obtain a high yield although the content of apatite in the ore is low.

CLAIMS


1. A froth flotation process for the enrichment of a calcium phosphate-containing mineral from an ore also containing calcium carbonate, characterized in that the process is performed in the presence, as a collector, of a derivative of aspartic acid of the formula



where R^{I} is a hydrophobic group containing a hydrocarbon group of 6-24 carbon atoms; R^{II} is an alkyl group with 1-7 carbon atoms or a group of the formula $(\text{B})_y\text{H}$, in which B is an alkyleneoxy group with 2-4 carbon atoms and y is a number from 1 to 10; and M is a group selected from the group consisting of a cation or hydrogen.

2. A froth flotation process in accordance with claim 1, characterized in that R^{I} is a glycidyl ether group of the formula $\text{CH}_2\text{CH}(\text{OH})\text{CH}_2\text{O}(\text{A}_1)_{x_1}\text{R}_1$, in which R_1 is a hydrocarbon group with 8-24 carbon atoms, A_1 is an alkyleneoxy group with 2-4 carbon atoms and x_1 is a number from 0 to 10; a hydroxyl group of the formula $\text{CH}_2\text{CH}(\text{OH})\text{R}_2$, in which R_2 is a hydrocarbon group with 6-22 carbon atoms; a propylene ether group of the formula $\text{C}_3\text{H}_6\text{O}(\text{A}_3)_{x_3}\text{R}_3$, in which R_3 is a hydrocarbon group with 8-24 carbon atoms, A_3 is an alkyleneoxy group with 2-4 carbon atoms and x_3 is a number from 0-10; or of the formula R_4 , where R_4 is a hydrocarbon group containing 8-24 carbon atoms.


3. A froth flotation process according to claim 2, characterized in that the derivative is selected from the group consisting of




where R_1 , A_1 , x_1 have the same meanings as in claim 2,



where R_2 has the same meaning as in claim 2,



where R_3 , A_3 and x_3 have the same meanings as in claim 2, and




where R_3 , A_3 and x_3 have the same meanings as in claim 2, or a mixture of two or more of the derivatives of formula II, III, IV or V.

4. A froth flotation process according to claim 2 or 3, characterized in that A_1 and A_3 is ethyleneoxy and x_1 and x_3 is a number from 1-4.

5. A froth flotation process according to claim 1 or 2, characterized in that R^{II} is methyl, hydroxyethyl or hydroxypropyl.


6. A froth flotation process according to any one of claims 1-5, characterized in that the derivative is present in an amount of 10-1500 grams per ton of the ore.

7. A froth flotation process according to any one of claims 1-6, characterized in that the process is performed in the presence of a polar co-collector of the formula




5 in which R^{III} is a hydrocarbon group with 8-22 carbon atoms, A is an oxyalkylene group having 2-4 carbon atoms and p is a number from 1-6,

or of the formula



15 in which R^{IV} is an aliphatic group having 7-21 carbon atoms, A is an alkyleneoxy group having 2-4 carbon atoms, q is a number from 0-6, and Y is an alkyl group having 1-4 carbon atoms or hydrogen, provided that Y cannot be hydrogen when q is zero.

8. A derivative of aspartic acid, characterized in that it has the formula




25 where R^I is a hydrophobic group containing a hydrocarbon group of 6-24 carbon atoms; R^{II} is an alkyl group with 1-7 carbons atoms or a group of the formula $(B)_yH$, in which B is an alkyleneoxy group with 2-4 carbon atoms and y is a number from 1 to 10; and M is a group selected from the group consisting of a cation or hydrogen.

9. A derivative according to claim 8, characterized in that R^I is a glycidylether group of the formula


30 $CH_2CH(OH)CH_2O(A_1)_{x1}R_1$, in which R_1 is a hydrocarbon group with 8-24 carbon atoms, A_1 is an alkyleneoxy group with 2-4 carbon atoms and $x1$ is a number from 0 to 10; a hydroxyl group of the formula $CH_2CH(OH)R_2$, in which R_2 is a hydrocarbon group

with 6-22 carbon atoms; a propylene ether group of the formula $C_3H_5O(A_3)_{x3}R_3$, in which R_3 is a hydrocarbon group with 8-24 carbon atoms, A_3 is an alkyleneoxy group with 2-4 carbon atoms and $x3$ is a number from 0-10, or of the formula R_4 , where R_4 is a hydrocarbon group containing 8-24 carbon atoms.


10. A derivative according to claim 9, characterized in that it is selected from the group consisting of




where R_1 , A_1 $x1$ have the same meanings as in claim 2,



where R_2 has the same meaning as in claim 2,




where R_3 , A_3 and $x3$ have the same meanings as in claim 2, and




where R_3 , A_3 and $x3$ have the same meanings as in claim 2, or a mixture of two or more of the derivatives of formula II, III, IV or V.

11. A method of producing a derivative according to claim 9, characterized in that maleic acid or a salt thereof is reacted under alkaline conditions with

a) a primary amine of the formula $R^{II}NH_2$, where R^{II} has the meaning mentioned above, followed by reacting the intermediate obtained with a glycidylether of the formula



where R_1 , x_1 and A_1 have the meanings mentioned above, an epoxide of the formula




where R_2 has the meaning mentioned above, or a halide compound of the formula $HalR_4$, where Hal is a halide and R_4 has the meaning above; or


b) with a primary amine of the formula $R^I NH_2$, where R^I has the meaning mentioned above, followed by reacting the intermediate obtained with a halide compound of the formula $HalR^{II}$, where Hal is a halide and R^{II} has the meaning mentioned above.

12. A method according to claim 11, characterized in that

i) the disodium salt of maleic acid is reacted with N-methylamine and the obtained (N-methyl)aspartate disodium salt is further reacted with a compound of the formula




where R_1 , A_1 and x_1 have the same meanings as in claim 11 to an aspartate of the formula II, or with a compound of the formula




where R_2 has the same meaning as in claim 2, to obtain an aspartate of the formula III, or

ii) the monosodium salt of maleic acid is reacted with an ether amine of the formula



where R_3 , A_3 and x_3 have the meanings mentioned in claim 11 to
5 obtain an intermediate of the formula




which intermediate is further reacted with $Cl(CH_2CH_2O)H$ or
10 CH_3Cl and with $NaOH$ to obtain a derivative of formula IV and V, respectively.

11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
66
67
68
69
70
71
72
73
74
75
76
77
78
79
80
81
82
83
84
85
86
87
88
89
90
91
92
93
94
95
96
97
98
99
100
101
102
103
104
105
106
107
108
109
110
111
112
113
114
115
116
117
118
119
120
121
122
123
124
125
126
127
128
129
130
131
132
133
134
135
136
137
138
139
140
141
142
143
144
145
146
147
148
149
150
151
152
153
154
155
156
157
158
159
160
161
162
163
164
165
166
167
168
169
170
171
172
173
174
175
176
177
178
179
180
181
182
183
184
185
186
187
188
189
190
191
192
193
194
195
196
197
198
199
200
201
202
203
204
205
206
207
208
209
210
211
212
213
214
215
216
217
218
219
220
221
222
223
224
225
226
227
228
229
230
231
232
233
234
235
236
237
238
239
240
241
242
243
244
245
246
247
248
249
250
251
252
253
254
255
256
257
258
259
260
261
262
263
264
265
266
267
268
269
270
271
272
273
274
275
276
277
278
279
280
281
282
283
284
285
286
287
288
289
290
291
292
293
294
295
296
297
298
299
300
301
302
303
304
305
306
307
308
309
310
311
312
313
314
315
316
317
318
319
320
321
322
323
324
325
326
327
328
329
330
331
332
333
334
335
336
337
338
339
340
341
342
343
344
345
346
347
348
349
350
351
352
353
354
355
356
357
358
359
360
361
362
363
364
365
366
367
368
369
370
371
372
373
374
375
376
377
378
379
380
381
382
383
384
385
386
387
388
389
390
391
392
393
394
395
396
397
398
399
400
401
402
403
404
405
406
407
408
409
410
411
412
413
414
415
416
417
418
419
420
421
422
423
424
425
426
427
428
429
430
431
432
433
434
435
436
437
438
439
440
441
442
443
444
445
446
447
448
449
450
451
452
453
454
455
456
457
458
459
460
461
462
463
464
465
466
467
468
469
470
471
472
473
474
475
476
477
478
479
480
481
482
483
484
485
486
487
488
489
490
491
492
493
494
495
496
497
498
499
500
501
502
503
504
505
506
507
508
509
510
511
512
513
514
515
516
517
518
519
520
521
522
523
524
525
526
527
528
529
530
531
532
533
534
535
536
537
538
539
540
541
542
543
544
545
546
547
548
549
550
551
552
553
554
555
556
557
558
559
560
561
562
563
564
565
566
567
568
569
570
571
572
573
574
575
576
577
578
579
580
581
582
583
584
585
586
587
588
589
590
591
592
593
594
595
596
597
598
599
600
601
602
603
604
605
606
607
608
609
610
611
612
613
614
615
616
617
618
619
620
621
622
623
624
625
626
627
628
629
630
631
632
633
634
635
636
637
638
639
640
641
642
643
644
645
646
647
648
649
650
651
652
653
654
655
656
657
658
659
660
661
662
663
664
665
666
667
668
669
670
671
672
673
674
675
676
677
678
679
680
681
682
683
684
685
686
687
688
689
690
691
692
693
694
695
696
697
698
699
700
701
702
703
704
705
706
707
708
709
710
711
712
713
714
715
716
717
718
719
720
721
722
723
724
725
726
727
728
729
730
731
732
733
734
735
736
737
738
739
740
741
742
743
744
745
746
747
748
749
750
751
752
753
754
755
756
757
758
759
760
761
762
763
764
765
766
767
768
769
770
771
772
773
774
775
776
777
778
779
780
781
782
783
784
785
786
787
788
789
790
791
792
793
794
795
796
797
798
799
800
801
802
803
804
805
806
807
808
809
810
811
812
813
814
815
816
817
818
819
820
821
822
823
824
825
826
827
828
829
830
831
832
833
834
835
836
837
838
839
840
841
842
843
844
845
846
847
848
849
850
851
852
853
854
855
856
857
858
859
860
861
862
863
864
865
866
867
868
869
870
871
872
873
874
875
876
877
878
879
880
881
882
883
884
885
886
887
888
889
890
891
892
893
894
895
896
897
898
899
900
901
902
903
904
905
906
907
908
909
910
911
912
913
914
915
916
917
918
919
920
921
922
923
924
925
926
927
928
929
930
931
932
933
934
935
936
937
938
939
940
941
942
943
944
945
946
947
948
949
950
951
952
953
954
955
956
957
958
959
960
961
962
963
964
965
966
967
968
969
970
971
972
973
974
975
976
977
978
979
980
981
982
983
984
985
986
987
988
989
990
991
992
993
994
995
996
997
998
999
1000
1001
1002
1003
1004
1005
1006
1007
1008
1009
1010
1011
1012
1013
1014
1015
1016
1017
1018
1019
1020
1021
1022
1023
1024
1025
1026
1027
1028
1029
1030
1031
1032
1033
1034
1035
1036
1037
1038
1039
1040
1041
1042
1043
1044
1045
1046
1047
1048
1049
1050
1051
1052
1053
1054
1055
1056
1057
1058
1059
1060
1061
1062
1063
1064
1065
1066
1067
1068
1069
1070
1071
1072
1073
1074
1075
1076
1077
1078
1079
1080
1081
1082
1083
1084
1085
1086
1087
1088
1089
1090
1091
1092
1093
1094
1095
1096
1097
1098
1099
1100
1101
1102
1103
1104
1105
1106
1107
1108
1109
1110
1111
1112
1113
1114
1115
1116
1117
1118
1119
1120
1121
1122
1123
1124
1125
1126
1127
1128
1129
1130
1131
1132
1133
1134
1135
1136
1137
1138
1139
1140
1141
1142
1143
1144
1145
1146
1147
1148
1149
1150
1151
1152
1153
1154
1155
1156
1157
1158
1159
1160
1161
1162
1163
1164
1165
1166
1167
1168
1169
1170
1171
1172
1173
1174
1175
1176
1177
1178
1179
1180
1181
1182
1183
1184
1185
1186
1187
1188
1189
1190
1191
1192
1193
1194
1195
1196
1197
1198
1199
1200
1201
1202
1203
1204
1205
1206
1207
1208
1209
1210
1211
1212
1213
1214
1215
1216
1217
1218
1219
1220
1221
1222
1223
1224
1225
1226
1227
1228
1229
1230
1231
1232
1233
1234
1235
1236
1237
1238
1239
1240
1241
1242
1243
1244
1245
1246
1247
1248
1249
1250
1251
1252
1253
1254
1255
1256
1257
1258
1259
1260
1261
1262
1263
1264
1265
1266
1267
1268
1269
1270
1271
1272
1273
1274
1275
1276
1277
1278
1279
1280
1281
1282
1283
1284
1285
1286
1287
1288
1289
1290
1291
1292
1293
1294
1295
1296
1297
1298
1299
1300
1301
1302
1303
1304
1305
1306
1307
1308
1309
1310
1311
1312
1313
1314
1315
1316
1317
1318
1319
1320
1321
1322
1323
1324
1325
1326
1327
1328
1329
1330
1331
1332
1333
1334
1335
1336
1337
1338
1339
1340
1341
1342
1343
1344
1345
1346
1347
1348
1349
1350
1351
1352
1353
1354
1355
1356
1357
1358
1359
1360
1361
1362
1363
1364
1365
1366
1367
1368
1369
1370
1371
1372
1373
1374
1375
1376
1377
1378
1379
1380
1381
1382
1383
1384
1385
1386
1387
1388
1389
1390
1391
1392
1393
1394
1395
1396
1397
1398
1399
1400
1401
1402
1403
1404
1405
1406
1407
1408
1409
1410
1411
1412
1413
1414
1415
1416
1417
1418
1419
1420
1421
1422
1423
1424
1425
1426
1427
1428
1429
1430
1431
1432
1433
1434
1435
1436
1437
1438
1439
1440
1441
1442
1443
1444
1445
1446
1447
1448
1449
1450
1451
1452
1453
1454
1455
1456
1457
1458
1459
1460
1461
1462
1463
1464
1465
1466
1467
1468
1469
1470
1471
1472
1473
1474
1475
1476
1477
1478
1479
1480
1481
1482
1483
1484
1485
1486
1487
1488
1489
1490
1491
1492
1493
1494
1495
1496
1497
1498
1499
1500
1501
1502
1503
1504
1505
1506
1507
1508
1509
1510
1511
1512
1513
1514
1515
1516
1517
1518
1519
1520
1521
1522
1523
1524
1525
1526
1527
1528
1529
1530
1531
1532
1533
1534
1535
1536
1537
1538
1539
1540
1541
1542
1543
1544
1545
1546
1547
1548
1549
1550
1551
1552
1553
1554
1555
1556
1557
1558
1559
1560
1561
1562
1563
1564
1565
1566
1567
1568
1569
1570
1571
1572
1573
1574
1575
1576
1577
1578
1579
1580
1581
1582
1583
1584
1585
1586
1587
1588
1589
1590
1591
1592
1593
1594
1595
1596
1597
1598
1599
1600
1601
1602
1603
1604
1605
1606
1607
1608
1609
1610
1611
1612
1613
1614
1615
1616
1617
1618
1619
1620
1621
1622
1623
1624
1625
1626
1627
1628
1629
1630
1631
1632
1633
1634
1635
1636
1637
1638
1639
1640
1641
1642
1643
1644
1645
1646
1647
1648
1649
1650
1651
1652
1653
1654
1655
1656
1657
1658
1659
1660
1661
1662
1663
1664
1665
1666
1667
1668
1669
1670
1671
1672
1673
1674
1675
1676
1677
1678
1679
1680
1681
1682
1683
1684
1685
1686
1687
1688
1689
1690
1691
1692
1693
1694
1695
1696
1697
1698
1699
1700
1701
1702
1703
1704
1705
1706
1707
1708
1709
1710
1711
1712
1713
1714
1715
1716
1717
1718
1719
1720
1721
1722
1723
1724
1725
1726
1727
1728
1729
1730
1731
1732
1733
1734
1735
1736
1737
1738
1739
1740
1741
1742
1743
1744
1745
1746
1747
1748
1749
1750
1751
1752
1753
1754
1755
1756
1757
1758
1759
1760
1761
1762
1763
1764
1765
1766
1767
1768
1769
1770
1771
1772
1773
1774
1775
1776
1777
1778
1779
1780
1781
1782
1783
1784
1785
1786
1787
1788
1789
1790
1791
1792
1793
1794
1795
1796
1797
1798
1799
1800
1801
1802
1803
1804
1805
1806
1807
1808
1809
1810
1811
1812
1813
1814
1815
1816
1817
1818
1819
1820
1821
1822
1823
1824
1825
1826
1827
1828
1829
1830
1831
1832
1833
1834
1835
1836
1837
1838
1839
1840
1841
1842
1843
1844
1845
1846
1847
1848
1849
1850
1851
1852
1853
1854
1855
1856
1857
1858
1859
1860
1861
1862
1863
1864
1865
1866
1867
1868
1869
1870
1871
1872
1873
1874
1875
1876
1877
1878
1879
1880
1881
1882
1883
1884
1885
1886
1887
1888
1889
1890
1891
1892
1893
1894
1895
1896
1897
1898
1899
1900
1901
1902
1903
1904
1905
1906
1907
1908
1909
1910
1911
1912
1913
1914
1915
1916
1917
1918
1919
1920
1921
1922
1923
1924
1925
1926
1927
1928
1929
1930
1931
1932
1933
1934
1935
1936
1937
1938
1939
1940
1941
1942
1943
1944
1945
1946
1947
1948
1949
1950
1951
1952
1953
1954
1955
1956
1957
1958
1959
1960
1961
1962
1963
1964
1965
1966
1967
1968
1969
1970
1971
1972
1973
1974
1975
1976
1977
1978
1979
1980
1981
1982
1983
1984
1985
1986
1987
1988
1989
1990
1991
1992
1993
1994
1995
1996
1997
1998
1999
2000
2001
2002
2003
2004
2005
2006
2007
2008
2009
2010
2011
2012
2013
2014
2015
2016
2017
2018
2019
2020
2021
2022
2023
2024
2025
2026
2027
2028
2029
2030
2031
2032
2033
2034
2035
2036
2037
2038
2039
2040
2041
2042
2043
2044
2045
2046
2047
2048
2049
2050
2051
2052
2053
2054
2055
2056
2057
2058
2059
2060
2061
2062
2063
2064
2065
2066
2067
2068
2069
2070
2071
2072
2073
2074
2075
2076
2077
2078
2079
2080
2081
2082
2083
2084
2085
2086
2087
2088
2089
2090
2091
2092
2093
2094
2095
2096
2097
2098
2099
2100
2101
2102
2103
2104
2105
2106
2107
2108
2109
2110
2111
2112
2113
2114
2115
2116
2117
2118
2119
2120
2121
2122
2123
2124
2125
2126
2127
2128
2129
2130
2131
2132
2133
2134
2135
2136
2137
2138
2139
2140
2141
2142
2143
2144
2145
2146
2147
2148
2149
2150
2151
2152
2153
2154
2155
2156
2157
2158
2159
2160
2161
2162
2163
2164
2165
2166
2167

ABSTRACT

A derivative of aspartic acid is used as a collector for a phosphate containing mineral, such as apatite, in a froth flotation process. According to the invention the collector has a high selectivity for phosphate containing minerals even in the presence of carbonate minerals, such as calcite.

The derivative has the formula



where R^{I} is a hydrophobic group containing a hydrocarbon group of 6-24 carbon atoms; R^{II} is an alkyl group with 1-7 carbon atoms or a group of the formula $(\text{B})_y\text{H}$, in which B is an alkyleneoxy group with 2-4 carbon atoms and y is a number from 1 to 10; and M is a group selected from the group consisting of a cation or hydrogen.

**This Page is Inserted by IFW Indexing and Scanning
Operations and is not part of the Official Record**

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☐ **BLACK BORDERS**
- ☐ **IMAGE CUT OFF AT TOP, BOTTOM OR SIDES**
- ☐ **FADED TEXT OR DRAWING**
- ☐ **BLURRED OR ILLEGIBLE TEXT OR DRAWING**
- ☐ **SKEWED/SLANTED IMAGES**
- ☐ **COLOR OR BLACK AND WHITE PHOTOGRAPHS**
- ☐ **GRAY SCALE DOCUMENTS**
- ☐ **LINES OR MARKS ON ORIGINAL DOCUMENT**
- ☐ **REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY**
- ☐ **OTHER:** _____

IMAGES ARE BEST AVAILABLE COPY.

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.